



Early View

Research letter

Pre-pubertal smoke exposure of fathers and increased risk of offspring asthma: a possible transgenerational effect

Jiacheng Liu, Gayan Bowatte, Jonathan Pham, Jennifer L. Perret, John W. Holloway, Adrian J. Lowe, John A. Burgess, Cecilie Svanes, Paul Thomas, Melissa A. Russell, Bircan Erbas, Caroline J. Lodge, David Martino, Gita D. Mishra, Michael J. Abramson, Eugene H. Walters, Shyamali C. Dharmage, Dinh S. Bui

Please cite this article as: Liu J, Bowatte G, Pham J, *et al.* Pre-pubertal smoke exposure of fathers and increased risk of offspring asthma: a possible transgenerational effect. *Eur Respir J* 2022; in press (<https://doi.org/10.1183/13993003.00257-2022>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Title page

Title

Pre-pubertal smoke exposure of fathers and increased risk of offspring asthma: a possible transgenerational effect

Authors' names

Jiacheng Liu¹, Gayan Bowatte^{1,2}, Jonathan Pham^{1,3}, Jennifer L. Perret^{1,4}, John W. Holloway⁵, Adrian J. Lowe¹, John A. Burgess¹, Cecilie Svanes⁶, Paul Thomas⁷, Melissa A. Russell¹, Bircan Erbas⁸, Caroline J. Lodge¹, David Martino⁹, Gita D. Mishra¹⁰, Michael J. Abramson¹¹, Eugene H. Walters^{1,12}, Shyamali C. Dharmage^{1*}, Dinh S. Bui^{1*}.

* Joint senior authors

Authors' affiliations

1. Allergy and Lung Health Unit, Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, VIC, Australia.
2. Department of Basic Sciences, Faculty of Allied Health Sciences, University of Peradeniya, Peradeniya, Sri Lanka.
3. Allergy, Asthma & Clinical Immunology Service, Department of Respiratory Medicine, The Alfred Hospital, Melbourne, VIC, Australia.
4. Institute for Breathing and Sleep, Heidelberg, VIC, Australia.
5. Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, United Kingdom; NIHR Southampton Biomedical Research Centre, University

Hospital Southampton, Southampton, United Kingdom.

6. Centre for International Health, Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway; Department of Occupational Medicine, Haukeland University Hospital, Bergen, Norway.
7. Prince of Wales' Clinical School, Faculty of Medicine, University of New South Wales, Sydney, NSW, Australia.
8. School of Psychology and Public Health, La Trobe University, Melbourne, VIC, Australia.
9. Wal-yan Respiratory Research Centre, Telethon Kids Institute, The University of Western Australia, Perth, WA, Australia.
10. Center for Longitudinal and Lifecourse Research, School of Public Health, The University of Queensland, Brisbane, QLD, Australia.
11. School of Public Health & Preventive Medicine, Monash University, Melbourne, VIC, Australia.
12. School of Medicine, University of Tasmania, Hobart, TAS, Australia.

Corresponding author: Professor Shyamali C Dharmage, Allergy and Lung Health Unit, Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Carlton, VIC 3053, Australia

Email: s.dharmage@unimelb.edu.au

All source(s) of funding

There is no funding to report for this submission.

Conflicts of interest

M.J.A holds investigator initiated grants from Pfizer, Boehringer-Ingelheim, Sanofi and GSK for unrelated research. He has undertaken an unrelated consultancy and received assistance with conference attendance from Sanofi. He has also received a speaker's fee from GSK. The rest of the authors declare that they have no conflicts of interest.

Key words

Asthma, offspring, paternal, passive smoking, transgenerational effect

Take-Home Message

The risk of non-allergic childhood asthma in offspring may increase if their fathers were exposed to pre-pubertal passive smoke.

Acknowledgements

This study is based on the data collected within the Tasmanian Longitudinal Health Study (TAHS).

The authors gratefully acknowledge all the participants and research staff of the TAHS.

Author contributions

Study conceptualization and design: J.L, G.B, J.P, J.L.P, D.S.B, S.C.D; Data analysis and interpretation: J.L, G.B, J.P, D.S.B, S.C.D; Data collection: G.B, J.P, J.L.P, J.W.H, A.J.L, J.A.B, C.S, P.T, M.A.R, B.E, C.J.L, D.M, G.D.M, M.J.A, E.H.W, D.S.B, S.C.D; Manuscript preparation: J.L, D.S.B, S.C.D; Manuscript modification: G.B, J.P, J.L.P, J.W.H, A.J.L, J.A.B, C.S, P.T, M.A.R, B.E, C.J.L, D.M, G.D.M, M.J.A, E.H.W. All authors critically reviewed the manuscript and approved the

final version for submission.

Human ethics approval declaration

This study was based on the Tasmanian Longitudinal Health Study, which was approved by Human Ethics Review Committees at the Universities of Melbourne (approval number: 040375), Tasmania (040375.1), New South Wales (08094), Alfred Hospital (1118/04), and Royal Brisbane and Women's Hospital Health Service District (2006/037).

Word count: 1392

Tables: 1; References: 15

Pre-pubertal smoke exposure of fathers and increased risk of offspring asthma: a possible transgenerational effect

To the Editors:

Asthma is a major non-communicable disease in children [1]. Pre- and post-natal exposure to tobacco smoke are major risk factors for childhood asthma [1, 2]. While there is evidence that mothers' intrauterine exposure to second-hand smoke is associated with asthma in the offspring [3, 4], there is also increasing concern that fathers who start smoking before completing puberty may elevate the risk of asthma in their offspring [4, 5]. The suggestion is that this may be as a result of epigenetic changes to sperm precursor (stem) cells during gonadal maturation [4, 5]. However, this is rather speculative, and as yet little is actually known about whether fathers' passive smoke exposure throughout childhood to puberty is indeed associated with increased asthma risk in their offspring.

We investigated this potential association using data from 1,689 father-offspring pairs in the Tasmanian Longitudinal Health Study (TAHS). Offspring in this study were the probands in TAHS [6], where their childhood asthma and hayfever status had been reported by their parents at age 7 years, in 1968. Offspring asthma was defined as more than one attack of asthma/wheezy breathing by this age of 7 years, and was classified as allergic asthma if they had both asthma and hayfever during the same period. Fathers of the probands reported their personal lifetime active smoking history and also pre-pubertal passive smoke exposure (between 0 and 15 years) from their own parents (i.e., offspring paternal grandparents) in the 2010 follow-up survey. We used logistic regression for offspring asthma at age 7 years (asthma vs. no asthma), and multinomial logistic regression for offspring asthma phenotypes (non-allergic asthma vs. no asthma; allergic asthma vs. no asthma). Analyses were adjusted for fathers' educational levels, and prior asthma history. These potential confounders were identified by directed acyclic graph (DAG) models. The analyses were additionally adjusted and stratified for fathers' lifetime active smoking history. In addition, we investigated potential interaction between fathers' pre-pubertal passive smoke exposure and lifetime active smoking, using a likelihood ratio test. We also combined fathers' pre-pubertal passive smoke exposure (yes/no) and fathers' lifetime active smoking history (yes/no), and categorised them into four groups (as: no/no [reference], yes/no, no/yes, yes/yes) and fitted this new variable in the analyses.

Fathers' passive smoke exposure before 15 years of age was associated with increased odds of offspring non-allergic asthma by 7 years of age (adjusted multinomial odds ratio [aMOR]: 1.59; 95% confidence interval [CI]: 1.09 – 2.32) (Adjusted model 1, Table 1). After further stratifying by fathers' lifetime active smoking history, the association between fathers' passive smoke exposure and offspring non-allergic asthma was stronger in ever-smoked fathers (n = 985; aMOR: 1.72; 95% CI: 1.02 – 2.92) than in fully non-smoking fathers (n = 601; aMOR: 1.39; 95% CI: 0.78 – 2.46). Further interaction analyses suggested a significant association of combined fathers' exposure to passive smoke before completing puberty and active smoking with offspring childhood non-allergic asthma (aMOR: 1.68; 95% CI: 1.02 – 2.79). There were increased odds of offspring childhood non-allergic asthma for fathers' exposure to pre-pubertal passive smoke only (aMOR: 1.38; 95% CI: 0.78 – 2.43), but not fathers' lifetime active smoking only (aMOR: 0.99; 95% CI: 0.51 – 1.94). Notably, the observed associations with offspring overall asthma (adjusted odds ratio [aOR]: 1.23; 95% CI: 0.92 – 1.65) or allergic asthma (aMOR: 0.84; 95% CI: 0.54 – 1.30) were not statistically significant at the 5% threshold.

Our findings suggest that when boys are passively exposed to their parent(s)' tobacco smoke before the age of 15 years, their offspring have increased risk of non-allergic childhood asthma, but not allergic asthma. These findings provide new insights and stronger evidence for the transgenerational effects caused by paternal passive smoke exposure on their offspring health. In comparison to our study that investigates the impact of fathers' pre-pubertal passive smoke exposure, one study reported an association between fathers' active smoking before the age of 15 years and their offspring's risk of non-allergic childhood asthma [5]. This would support our findings, given the similar timing of exposure, albeit likely differing quantity and constituents of smoke exposure. However, our findings are novel as they provide the first evidence on transgenerational transmission of an adverse impact of pre-pubertal passive smoke exposure in fathers.

Epigenetics is a potential mechanistic link between paternal pre-pubertal smoking and offspring elevated asthma risk [4, 5], via changes in sperm stem cell including DNA methylation [5] and altered miRNA profiles [4]. Further, offspring prenatal passive smoke exposure may modify their responses to nicotine metabolites and other smoking related substances [7-9], and adversely affect

lung functional development and their immunity in later life [4, 7, 10]. For example, one rat study demonstrated potentially heritable nicotine-induced DNA methylation [11], and increased CG methylation levels of immune genes in human offspring by 6 years of age were attributed to prenatal paternal smoking and related to childhood asthma [10]. Therefore, the detrimental effect of fathers' early life exposure to any type of tobacco smoke may be epigenetically transmitted to the next generation. Any differences in this between passive smoke exposure and active smoking requires further investigations.

Our findings also suggest that abstinence from active smoking by father could potentially attenuate offspring non-allergic asthma risk imposed by the fathers' pre-pubertal passive smoke exposure, and this needs to be followed up. This finding could be supported by the hypothesis that although epigenetic programming is in part heritable, there is a developmental reprogramming process, in which DNA methylation and histone modifications may be reset or erased at every generation to different degrees [7]. In other words, there might be an adaptive response or maintenance mechanism that prevents further damage [5]. There is evidence that initial smoking-related DNA methylation changes can partially recover following smoking cessation [12]. One murine model showed that the perinatal nicotine-induced asthmatic lung phenotype tended to be less prominent in successive generations that were no longer exposed, but was exacerbated after re-exposure during the second pregnancy [11]. These findings together with ours suggest that fathers' active smoking may exacerbate the impact of their passive smoking before the age of 15 years on offspring asthma risk. Pre-pubertal passive smoking exposed fathers without the insult from active smoking may lead to an attenuated risk on offspring.

Asthma is a heterogeneous disease [13]. In the modern era of precision medicine, understanding risk factors for different asthma phenotypes is important, and potentially for treatment responsiveness [14]. Our data showed that paternal smoke exposure before the age of 15 years is a major risk factor for non-allergic asthma, shedding light on the potential epigenetic and pathophysiological mechanisms of disease. Furthermore, non-allergic asthma may be less responsive to inhaled corticosteroids and so more difficult to treat [15]. Therefore, public health interventions to reduce early-life smoke exposure could help prevent difficult-to-treat asthma in offspring. Moreover, our findings may be valuable to direct researchers to conduct further

investigations to identify specific mechanisms and propose a new paradigm of non-allergic asthma treatments from epigenetic level in the future, such as treatments to reduce epigenetic effects of smoke exposure. Lastly, our results revealed some limited evidence for an increased risk of overall asthma (point estimate increased by 23% but CIs included the null value of 1). Power was possibly the issue. The association for allergic asthma has a wide confidence interval, thus our result was inconclusive to determine whether passive smoke exposure of fathers influenced risk of allergic asthma in their offspring. More studies with larger sample size are required.

TAHS is a population-based birth cohort study which has collected longitudinal data and also among proband relatives in multiple generations [6], which is a major strength. Nevertheless, we acknowledge that fathers' reports of their smoke exposure was potentially subject to misclassification. However, fathers' passive smoke exposure and smoking history were ascertained in 2010, which was far away from their report of offspring asthma in 1968; report of their smoke exposure was less likely to be influenced by offspring asthma status. Hence, any such misclassification was more likely to be non-differential, which would only bias the associations towards the null. We were unable to validate fathers' report of their smoke exposure, which is a limitation of this study.

In conclusion, this study suggests that the risk of non-allergic childhood asthma in offspring may increase if their fathers had been exposed to pre-pubertal passive smoke. Avoidance of smoking, in addition to the multitude of known health benefits, can also reduce asthma risk in offspring.

Table 1. Association between fathers' passive smoke exposure before the age of 15 years and offspring non-allergic asthma by the age of 7 years

Exposure	Offspring non-allergic asthma by the age of 7 years									
	Full sample (n = 1586)			Non-smoking fathers (n = 601)			Ever-smoked fathers (n = 985)			
Fathers' passive smoke exposure before the age of 15 years from their own parent(s)	Crude model	Adjusted model 1	Adjusted mode 2	Crude model	Adjusted model 1	Crude model	Adjusted model 1	Cases/total	MOR	MOR
	MOR	MOR	MOR	MOR	MOR	MOR	MOR	MOR	MOR	MOR
	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)
	Cases/total			Cases/total			Cases/total			
Not exposed (ref)	38/455	1	1	1	20/240	1	1	18/215	1	1
Exposed	144/1131	1.60	1.59	1.54	40/361	1.37	1.39	104/770	1.71	1.72
		(1.10, 2.33)*	(1.09,2.32)*	(1.05,2.26)*		(0.78,2.41)	(0.78,2.46)		(1.01,2.89)*	(1.02,2.92)*

* $P < 0.05$

Adjusted model 1: Adjustment for fathers' educational levels and fathers' prior asthma history

Adjusted model 2: Model 1 plus further adjustment for fathers' lifetime active smoking history

References

1. Harju M, Keski-Nisula L, Georgiadis L, et al. Parental smoking and cessation during pregnancy and the risk of childhood asthma. *BMC Public Health* 2016;16:428.
2. Tabuchi T, Fujiwara T, Nakayama T, et al. Maternal and paternal indoor or outdoor smoking and the risk of asthma in their children: a nationwide prospective birth cohort study. *Drug Alcohol Depend* 2015;147:103-108.
3. Magnus MC, Haberg SE, Karlstad O, et al. Grandmother's smoking when pregnant with the mother and asthma in the grandchild: the Norwegian Mother and Child Cohort Study. *Thorax* 2015;70(3):237-243.
4. Accordini S, Calciano L, Johannessen A, et al. A three-generation study on the association of tobacco smoking with asthma. *Int J Epidemiol* 2018;47(4):1106-1117.
5. Svanes C, Koplun J, Skulstad SM, et al. Father's environment before conception and asthma risk in his children: a multi-generation analysis of the Respiratory Health In Northern Europe study. *Int J Epidemiol* 2017;46(1):235-245.
6. Matheson MC, Abramson MJ, Allen K, et al. Cohort Profile: The Tasmanian Longitudinal Health STUDY (TAHS). *Int J Epidemiol* 2017;46(2):407-408i.
7. Krauss-Etschmann S, Meyer KF, Dehmel S, et al. Inter- and transgenerational epigenetic inheritance: evidence in asthma and COPD? *Clin Epigenetics* 2015;7:53.
8. Harlid S, Xu Z, Panduri V, et al. CpG sites associated with cigarette smoking: analysis of epigenome-wide data from the Sister Study. *Environ Health Perspect*. 2014;122(7):673-678.
9. Liu G, Lin CJ, Yates CR, et al. Metabolomic Analysis Identified Reduced Levels of Xenobiotics, Oxidative Stress, and Improved Vitamin Metabolism in Smokers Switched to Vuse Electronic Nicotine Delivery System. *Nicotine Tob Res*. 2021;23(7):1133-1142.
10. Wu CC, Hsu TY, Chang JC, et al. Paternal Tobacco Smoke Correlated to Offspring Asthma and Prenatal Epigenetic Programming. *Front Genet* 2019;10:471.
11. Liu J, Yu C, Doherty TM, Akbari O, et al. Perinatal nicotine exposure-induced transgenerational asthma: Effects of reexposure in F1 gestation. *FASEB J* 2020;34(9):11444-11459.

12. Tsaprouni LG, Yang TP, Bell J, et al. Cigarette smoking reduces DNA methylation levels at multiple genomic loci but the effect is partially reversible upon cessation. *Epigenetics* 2014;9(10):1382-1396.
13. Bui DS, Lodge CJ, Perret JL, et al. Trajectories of asthma and allergies from 7 years to 53 years and associations with lung function and extrapulmonary comorbidity profiles: a prospective cohort study. *Lancet Respir Med.* 2021;9(4):387-396.
14. Chau-Etchepare F, Hoerger JL, Kuhn BT, et al. Viruses and non-allergen environmental triggers in asthma. *J Investig Med.* 2019;67(7):1029-1041.
15. Marcon A, Marchetti P, Antó JM, et al; Atopy Modifies the Association Between Inhaled Corticosteroid Use and Lung Function Decline in Patients with Asthma. *J Allergy Clin Immunol Pract.* 2020;8(3):980-988.