



Early View

Correspondence

The association between SARS-CoV-2 RT-PCR cycle threshold and mortality in a community cohort

Rupert Waudby-West, Benjamin Parcell, Colin N A Palmer, Samira Bell, James D Chalmers, Moneeza K Siddiqui

Please cite this article as: Waudby-West R, Parcell B, Palmer CNA, *et al.* The association between SARS-CoV-2 RT-PCR cycle threshold and mortality in a community cohort. *Eur Respir J* 2021; in press (<https://doi.org/10.1183/13993003.00360-2021>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

The association between SARS-CoV-2 RT-PCR cycle threshold and mortality in a community cohort

Rupert Waudby-West¹, Benjamin Parcell² Colin N A Palmer¹, Samira Bell¹, James D Chalmers³, Moneeza K Siddiqui¹

1. Division of Population Health and Genomics, School of Medicine, University of Dundee, Dundee DD1 9SY

2. Department of Medical Microbiology Ninewells Hospital & Medical School, Dundee DD1 9SY

3. Division of Molecular and Clinical Medicine, School of Medicine, University of Dundee, DD1 9SY

Corresponding author:

Prof. James D. Chalmers
Division of Molecular & Clinical Medicine
School of Medicine
University of Dundee
Email: j.chalmers@dundee.ac.uk

Tweet: Lower Ct values from RT-PCR tests are associated with higher risk of mortality in SARS-CoV2 infections. Ct values <20 was associated with 2.20 times the hazards of mortality compared to values >30 [(95%CI: 1.28-3.76) adjusted for age, sex, comorbidities & hospitalisation

A diagnosis of severe acute respiratory virus coronavirus 2 (SARS-CoV-2) disease is widely made by the use of reverse transcription polymerase chain reaction (RT-PCR). When using RT-PCR, the infectivity of SARS-CoV-2 can be inferred from the threshold cycle (Ct) value [1]. It is best practice to confirm the validity of the standard curve using reference materials or in-house plasmid controls with known viral copy numbers [2]. As the Ct value represents the cycle number at which the signal breaches the threshold for positivity, a lower Ct value is indicative of a higher viral load. Although some studies suggest that viral load is associated with mortality and infectiousness [3,4], a systematic review has identified little difference in viral load between pre-symptomatic, asymptomatic and symptomatic patients [5]. As a result, the clinical relevance of viral load remains controversial, and it is not used in clinical practice [6]. Here, we report the relationship between the Ct value and all-cause mortality for people who tested positive for SARS-CoV-2 on a combined nasal and pharyngeal swab in the Tayside region of Scotland. This is a community cohort study

and includes the local population of the region, as well as symptomatic health and social care workers tested as part of a screening programme [7]. In order to obtain clinical characteristics and outcomes for those who tested positive, anonymised record linkage was conducted between routine healthcare datasets as described previously [8]. All positive PCR tests from the 12th of March until the 1st of May 2020 were included, and all deaths recorded by National Records Scotland until the 20th of May 2020. Approval for anonymised data linkage was granted by the local Data Protection Officer (Caldicott Guardian).

A total of 1337 individuals testing positive for SARS-CoV-2 were recorded in NHS Tayside. In the laboratory, nucleic acid was extracted and a one-step RT-PCR assay targeting the RdRP gene using an ABI7500 as previously described. For individuals with multiple positive tests, the Ct value from the first of these was included, and for multiple tests on the same day, the test with the lowest Ct value was used. In our cohort, the overall median Ct value of an individual's first positive test was 28.7 (IQR = 23.9–33.4). The median age of the cohort was 53 years (IQR = 40–72), and 61.9% were female. 202 individuals (15%) were hospitalised with COVID-19 disease (recorded with the ICD code for COVID-19: U07.1).

Total number of deaths in the cohort were 161 (12%), the majority of deaths (102) occurred in a non-hospital cohort. However, a larger proportion of those hospitalised died (29% compared to 9% who died without hospitalisation). This is likely due to the high incidence of infection and deaths in care homes for the elderly. Mean follow up available was 37 days (SD 14), and maximum follow up was 69 days. 15% of the cohort had hypertension, 4% had heart failure, 9% had ischaemic heart disease (IHD), 8% had chronic obstructive pulmonary disease (COPD), 5% had asthma, 8% had chronic kidney disease (CKD), and 10% had type 2 diabetes mellitus (T2DM).

A Cox proportional hazards model adjusting for relevant clinical factors was used to test the association between CT value and mortality. These were: age, sex, hypertension, heart failure, IHD, COPD, asthma, CKD, and T2DM. The model showed a significant association between initial Ct value and mortality (HR 0.96 [95% CI 0.93–0.99] $p = 0.0044$). Other independent predictors of mortality were age

(HR 1.08 [95% CI 1.06–1.09] $p < 0.001$), and male sex (HR 1.77 [95% CI 1.28–2.44] $p < 0.001$).

Ct values were then stratified into high (>30), moderate (20-30), and low (< 20), and a model was constructed adjusting for the same clinical factors. This showed that a low initial Ct value was associated with increased hazard of mortality compared to a high initial Ct value (HR 2.10 [95% CI 1.23-3.60] $p = 0.0069$), however the same relationship between moderate and high was not statistically significant (HR 1.37 [95% CI 0.96-1.94] $p = 0.08$). We performed a sensitivity analysis adjusting for hospitalisation status as a marker of disease severity; in this adjusted analysis a low initial Ct value increased hazards of mortality by 2.20 [(95%CI: 1.28-3.76) $p=0.0005$]. These results suggest that disease severity does not confound the association between Ct values and mortality.

Early identification of the risk that COVID-19 disease presents to an individual remains a challenge. Most clinical investigations for COVID-19 disease have been limited to hospitalised cohorts, here we are able to assess the value of Ct scores in the community. However, as this is a community cohort study, data from clinical investigations performed on hospitalised patients which help classify disease severity, are not available for the entire cohort. Here, we have demonstrated an independent relationship between the Ct value (indicative of viral load) of an individual's first positive SARS-CoV-2 PCR test, and overall mortality, in which a lower Ct value is associated with greater hazard of death. Ct values may allow for early risk stratification in patients with COVID-19. Further work is required to confirm this association and to assess the utility of including Ct values in early risk stratification.

References:

1. Public Health England. Understanding cycle threshold (Ct) in SARS-CoV-2 RT-PCR. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/926410/Understanding_Cycle_Threshold__Ct__in_SARS-CoV-2_RT-PCR_.pdf. Date last updated: October 28 2020. Date last accessed: January 17 2020.
2. Han MS, Byun JH, Cho Y, Rim JH. RT-PCR for SARS-CoV-2: quantitative versus qualitative. *Lancet Infect Dis*. 2020 May 20:S1473-3099(20)30424-2.
3. Pujadas E, Chaudhry F, McBride R, Richter F, Zhao S, Wajnberg A, Nadkarni G, Glicksberg BS, Houldsworth J, Cordon-Cardo C. SARS-CoV-2 viral load predicts COVID-19 mortality. *Lancet Respir Med*. 2020 Sep;8(9):e70
4. Singanayagam A, Patel M, Charlett A, Lopez Bernal J, Saliba V, Ellis J, Ladhani S, Zambon M, Gopal R. Duration of infectiousness and correlation with RT-PCR cycle threshold values in cases of COVID-19, England, January to May 2020. *Euro Surveill*. 2020 Aug;25(32):2001483.
5. Walsh KA, Jordan K, Clyne B, Rohde D, Drummond L, Byrne P, Ahern S, Carty PG, O'Brien KK, O'Murchu E, O'Neill M, Smith SM, Ryan M, Harrington P. SARS-CoV-2 detection, viral load and infectivity over the course of an infection. *J Infect*. 2020 Sep;81(3):357-371.
6. Kelleni MT. SARS CoV-2 viral load might not be the right predictor of COVID-19 mortality. *J Infect*. 2020 Aug 15:S0163-4453(20)30552-1.
7. Parcell BJ, Brechin K, Allstaff S, Park M, Third W, Bean S, Hind C, Farmer R, Chandler D, Chalmers JD. Drive-through testing for SARS-CoV-2 in symptomatic health and social care workers and household members: an observational cohort study. *Thorax*. 2020 Dec;75(12):1109-1111.
8. Siddiqui MK, Parcell BJ, Allstaff S, Palmer C, Chalmers JD, Bell S. Characteristics and outcomes of health and social care workers testing positive for SARS-CoV-2 in the Tayside region of Scotland. *Eur Respir J*. 2020 Sep 17;56(3):2002568.




Figure 1. **A.** Overall survival probability for individuals positive SARS-CoV-2 in Tayside, Scotland based on the cycle threshold (Ct) value of their first positive test. Crosses represent censored individuals. **B.** Estimated hazard ratio based on initial Ct value of individual in a model adjusting for age, sex, hypertension, heart failure, IHD, COPD, asthma, CKD, and T2DM.